

## CASE PROCESSING FOR ORGANS THAT WILL UNDERGO ISLET ISOLATION

### 1. PURPOSE

The purpose of this Standard Operating Procedure (SOP) is to outline procedures for processing pancreas and associated organs in conjunction with islet isolation for the nPOD Islet Isolation Pilot Program.

### 2. SCOPE

This SOP will be applied to all samples recovered through the nPOD Islet Isolation Pilot Program and designed to be used in conjunction with SOP IS-001(nPOD) "Human Pancreatic Islet Isolation For nPOD Islet Pilot Project."

### 3. RESPONSIBILITIES

3.1 Managers and supervisors are responsible for making sure that technicians are properly trained and equipment and facility are maintained in good working order.

3.2 Laboratory personnel are responsible for reading and understanding this SOP and related documents and to perform these tasks in accordance with the SOPs. They are responsible for following clinical laboratory and tissue banking best practices.

### 4. EQUIPMENT and MATERIALS

The materials, equipment, and forms listed in the following list are recommendations only and alternative products as suitable may be substituted for the site specific task or procedure.

- Kimberly-Clark Benchtop Protector, Fisher Scientific #15-235-101
- Leica Biosystem 3P L-absorb towel, Leica #3803240
- Double Edge Pathology Blades for Pathco Handle, Fisher Scientific #23-720-200
- Disposable Scalpel Handle with Blade 20/Pk, Fisher Scientific #5-0949-246
- Margin Marking Dye, Blue, American Master Tech #STD02204E
- Wooden Applicators with Cotton tips, Fisher Scientific #14-959-96B
- 32 oz. screw top polypropylene histology container, Fisher Scientific #22-026-315
- 52 oz. Plastic Container with Lid, Fisher Scientific #02-544-127
- Formalin, 1:10 dilution (buffered), Fisher Scientific #23-245-685
- Cassettes for embedding, blue, Lab Storage Systems #EC-0112

JDRF nPOD Standard Operating Procedure		
SOP Number and Version: 45.0	Supersedes: N/A	SPECIAL PROJECTS: ORGAN PROCUREMENT & PROCESSING CORE PROCEDURE
Original Effective Date: 08/27/2014		
Version Effective Date:		

Disposable Base Mold, Fisher Scientific #22-050-161  
 Microtubes with Silicon O-ring, VWR #89004-300  
 Tissue-Tek OCT Compound Media, Fisher Scientific #14-373-65  
 Ambion RNALater Solution, Ambion #AM7021  
 HyClone D-PBS 1X without Ca or Mg, Fisher Scientific #SH3002801  
 DMEM/F12 1:1 Modified, Fisher Scientific #SH30261.01  
 Anti-Anti, Antibiotic-Antimycotic 100X, Invitrogen #15240-062, aliquot 5ml and store at -20°C  
 Ammonium Chloride Solution, StemCell #7800  
 HyClone Fetal Bovine Serum, Fisher Scientific #SH30088031H, aliquot 50ml and store at -20°C  
 High-Performance centrifuge tubes (15, 50 ml), sterile, VWR, # 89039-666; #89039-65  
 Permanent marker (labeling OCT, etc), Fisher Scientific # 13383C  
 Kimberly-Clark Fluidshield Fog-Free Protective Mask, Fisher Scientific #18-806  
 Polyethylene Apron, Fisher Scientific # 19-181-529  
 Gloves  
 Ultra Clorox Germicidal Bleach  
 Ethanol  
 Biological Safety Hood  
 Centrifuge, room temperature  
 Organ/Tissue Scale  
 Dissecting Board, Fisher Scientific #36-114  
 Pathco Blade Handle, Fisher Scientific #NC9552049  
 Specimen Forceps, VWR #82027-438  
 Walter Stern Dissecting Scissors 310-045, LabSource 25876009  
 Drummond Pipet-Aid Filler/Dispenser, Fisher Scientific #13-681-15E  
 Pipettes, ThermoScientific Finnpiquette F2, 20-200 µl, Fisher Scientific #14-386-319  
 Pipettes, ThermoScientific Finnpiquette F2, 100-1000 µl Fisher Scientific # 13-386-320  
 ART Barrier Pipette tips 200ul, sterile, Thermo Scientific # 2279  
 ART Barrier Pipette tips 1000ul, sterile, Thermo Scientific # 2069  
 Biohazard Sharps Container  
 Styrofoam container  
 Liquid nitrogen

5. SAFETY

- 5.1. Use universal safety precautions when handling human samples and personal protective equipment (e.g. face mask with shield, gloves, lab coat or apron).
- 5.2. Follow chemical safety procedures and dispose of waste tissues according to UF EHS guidelines.
- 5.3. Handle sharps (e.g. scalpels, blood tubes) carefully and dispose properly.
- 5.4. Follow aseptic procedures throughout processing

<b>JDRF nPOD Standard Operating Procedure</b>		
SOP Number and Version: 45.0	Supersedes: N/A	SPECIAL PROJECTS: ORGAN PROCUREMENT & PROCESSING CORE PROCEDURE
Original Effective Date: 08/27/2014		
Version Effective Date:		
		Page 2 of 7

6. PROCEDURE

6.1. Preparation for sample processing

6.1.1. Case number assignment

6.1.1.1. Donors accepted as part of the nPOD Islet Isolation Pilot Project will be assigned case numbers in sequence with those at the University of Florida nPOD processing facility.

6.1.2. Aliquot labeling

6.1.2.1. Cassettes for paraffin embedding

6.1.2.1.1. Line #1: Case ID + Block #

6.1.2.1.2. Line #2: Sample type abbreviation

6.1.2.2. OCT cryomolds

6.1.2.2.1. Print legibly using a Sharpie permanent marker

6.1.2.2.2. Identify by Case ID, Block #, and sample type

6.1.2.3. Cryovials and Fresh Tissue

6.1.2.3.1. Line #1: Case ID + vial #

6.1.2.3.2. Line #2: Sample type abbreviation

6.1.2.3.3. Line #3: Aliquot type

6.1.2.3.4. Line #4: Barcode and unique aliquot number

6.1.3. Sterile media preparation

6.1.3.1. Remove and discard 50ml from 500ml RPMI stock solution.

6.1.3.2. Add 50ml FBS to the 450ml RPMI stock solution for a final concentration of 10% FBS.

6.1.3.3. Add 5ml 100X Antibiotic/antimycotic stock to the solution in 6.1.3.2.

6.1.3.4. Label the sterile media container with preparation date, additives, and preparer's initials. Store at 4°C for up to one month.

6.1.4. Liquid Nitrogen freezing bath

6.1.4.1. Pour liquid nitrogen into Styrofoam container and cover.

6.1.4.2. Place vial rack in the liquid nitrogen filled Styrofoam container and cover.

6.1.5. 10% NBF container

6.1.5.1. Pour 10% NBF into a 52 oz. plastic container and secure the lid.

6.1.6. Data Collection

6.1.6.1. Unpack shipment and record contents.

6.1.6.1.1. If any of the expected items were not received, notify the administrator on-call immediately.

6.1.6.2. Manually fill out the Case Worksheet form during the case processing.

6.1.6.2.1. After case has been processed scan, upload, and archive the completed worksheet. Manually enter the case information into the nPOD database.

6.2. Tissue Dissection

6.2.1.1. Receive one approximately 2.0 cm wide piece of pancreas from the PanHead/Body junction, and one approximately 2.0 cm wide piece of pancreas from

JDRF nPOD Standard Operating Procedure		
SOP Number and Version: 45.0	Supersedes: N/A	SPECIAL PROJECTS: ORGAN PROCUREMENT & PROCESSING CORE PROCEDURE
Original Effective Date: 08/27/2014		
Version Effective Date:		
		Page 3 of 7

the distal portion of the PanTail from islet isolation staff and move to hood, as cut by the Islet Lab Core. Make 5 blocks per 2.0 cm region: 2 paraffin and 2 OCT. The remaining piece divided into 2 cryovials, and 2 snap frozen.

- 6.2.1.1.1. Alternate paraffin and OCT as seen in Appendix 1.
- 6.2.1.1.2. Maintain the same tissue orientation for all blocks.
- 6.2.1.1.3. Paraffin blocks
  - 6.2.1.1.3.1. Place the tissue in the cassette.
  - 6.2.1.1.3.2. Attach and secure the top of the cassette.
  - 6.2.1.1.3.3. Place the cassette containing the tissue in the container of 10% NBF.
- 6.2.1.1.4. OCT blocks
  - 6.2.1.1.4.1. Place enough OCT media to cover the bottom surface of the cryomold.
  - 6.2.1.1.4.2. Place tissue on top of the media in the center of the mold and add OCT media until the tissue is covered.
  - 6.2.1.1.4.3. Place the filled cryomold on the top of the rack in the container with liquid nitrogen until frozen.
  - 6.2.1.1.4.4. Once frozen, move the OCT block to the polyurethane container with dry ice only.

6.2.2. Receive whole spleen and duodenum from islet isolation staff and move to hood.

6.2.3. Dissect the spleen into 2 regions.

- 6.2.3.1. 10cm x 20cm piece to be processed for cryovials, paraffin blocks, and OCT blocks.
- 6.2.3.2. The second piece is split into 5g aliquots, minced and transferred to 50ml conical tubes that contain 35ml of fresh complete sterile media. Secure the top and wrap with parafilm.
  - 6.2.3.2.1. The number of aliquots is determined by the number of investigators that will be receiving fresh spleen.

6.2.4. Process 10cm x 20 cm piece of spleen into snap frozen cryovials, paraffin blocks, and OCT blocks.

- 6.2.4.1. Snap frozen cryovials
  - 6.2.4.1.1. Without RNALater
    - 6.2.4.1.1.1. Mince 0.5 grams of tissue and place in vial.
    - 6.2.4.1.1.2. Secure the top of the vial and place in the vial rack in the container with liquid nitrogen.
  - 6.2.4.1.2. With RNALater
    - 6.2.4.1.2.1. Add 1ml RNALater to each vial
    - 6.2.4.1.2.2. Mince 0.5 grams of tissue and place in vial.
    - 6.2.4.1.2.3. Secure the top of the vial and allow the tissue and RNALater to equilibrate to room temperature for 30 minutes.
    - 6.2.4.1.2.4. Place the vial in the vial rack in the container with liquid nitrogen.

6.2.4.2. Paraffin blocks

<b>JDRF nPOD Standard Operating Procedure</b>		
SOP Number and Version: 45.0	Supersedes: N/A	SPECIAL PROJECTS: ORGAN PROCUREMENT & PROCESSING CORE PROCEDURE
Original Effective Date: 08/27/2014		
Version Effective Date:		Page 4 of 7


- 6.2.4.2.1. Place the tissue in the tissue cassette.
- 6.2.4.2.2. Attach and secure the top of the cassette.
- 6.2.4.2.3. Place the cassette containing the tissue in the container filled with 10% NBF.
- 6.2.4.3. OCT Blocks
  - 6.2.4.3.1. Place enough OCT media to cover the bottom surface of the cryomold.
  - 6.2.4.3.2. Place tissue on top of the media in the center of the mold and add OCT media until the tissue is covered.
  - 6.2.4.3.3. Place the filled cryomold on top of the rack in the container with liquid nitrogen until frozen.
- 6.2.5. In the hood, open the duodenum and gently scrape off mucus and ingesta.
- 6.2.6. Cut several 1" segments of mucosa.
  - 6.2.6.1. Fresh tissue for distribution
    - 6.2.6.1.1. Fill a conical tube to the 15ml mark with 1" segments of mucosa. Add fresh sterile media until the volume reaches 50ml.
  - 6.2.6.2. Snap frozen cryovials
    - 6.2.6.2.1. Without RNALater
      - 6.2.6.2.1.1. Mince a 1" strip of duodenum and place in vial.
      - 6.2.6.2.1.2. Secure the top of the vial and place the vial in the vial rack in the container with liquid nitrogen.
    - 6.2.6.2.2. With RNALater
      - 6.2.6.2.2.1. Add 1ml RNALater to vial
      - 6.2.6.2.2.2. Mince a 1" strip of duodenum and place in vial.
      - 6.2.6.2.2.3. Secure the top of the vial and allow the tissue and RNALater to equilibrate to room temperature for 30 minutes.
      - 6.2.6.2.2.4. Place the vial in the vial rack in the container with liquid nitrogen.
  - 6.2.6.3. Paraffin blocks
    - 6.2.6.3.1. Place a single 1" segment per cassette.
    - 6.2.6.3.2. Attach and secure the top of the cassette.
    - 6.2.6.3.3. Place the cassette containing the tissue in the container filled with 10% NBF.
  - 6.2.6.4. OCT blocks
    - 6.2.6.4.1. Place enough OCT media to cover the bottom surface of the cryomold.
    - 6.2.6.4.2. Place a 1" segment of duodenum on top of the media in the center of the mold and add OCT media until the tissue is covered.
    - 6.2.6.4.3. Place the filled cryomold on top of the rack in the container with liquid nitrogen.
- 6.2.7. In the hood, place the peri-pancreatic fat in the 32oz screw top polypropylene histology container. Fill the remaining volume of the container with fresh sterile media. Secure the top and wrap with parafilm.

<b>JDRF nPOD Standard Operating Procedure</b>		
SOP Number and Version: 45.0	Supersedes: N/A	SPECIAL PROJECTS: ORGAN PROCUREMENT & PROCESSING CORE PROCEDURE
Original Effective Date: 08/27/2014		
Version Effective Date:		Page 5 of 7

- 6.2.8.If thymus is received, dissect it into 2 equal sections.
- 6.2.8.1. The first section is to be cut into several pieces and transferred to 50ml conical tubes that contain 35ml of fresh complete sterile media. Secure the top and wrap with parafilm.
  - 6.2.8.2. The second section is to be cut in half and used to make 2 paraffin blocks and 2 OCT blocks.
  - 6.2.8.3. Paraffin blocks
    - 6.2.8.3.1. Place the tissue in the cassette.
    - 6.2.8.3.2. Attach and secure the top of the cassette.
    - 6.2.8.3.3. Place the cassette containing the tissue in the container filled with 10% NBF.
  - 6.2.8.4. OCT Blocks
    - 6.2.8.4.1. Place enough OCT media to cover the bottom surface of the cryomold.
    - 6.2.8.4.2. Place tissue on top of the media in the center of the mold and add OCT media until the tissue is covered.
    - 6.2.8.4.3. Place the filled cryomold on top of the rack in the container with liquid nitrogen.
- 6.2.9.After all tissue in the hood has been processed, centrifuge serum tubes at 1400 rpm for 10 minutes at room temperature.
- 6.2.9.1. If hemolysis is observed, record the degree (i.e. light or gross) and centrifuge for an additional 10 minutes at the same settings.
- 6.2.10. Aliquot the serum into labeled O-ring cryovials using volumes of 300ul or more per vial. Store at -80°C.
- 6.2.10.1. Set aside 3 aliquots for autoantibody and C-peptide analysis. Refer to SOP 85 C-Peptide Determination, and SOP 22 Autoantibody Screening Process.
- 6.2.11. All sample types that were frozen in the liquid nitrogen bath should be stored in a -80°C freezer until they are shipped to the OPPC Lab on dry ice.
- 6.2.12. The cassettes for paraffin processing are to be kept in 10% NBF for 24 hours and then processed according to the SOP 57.3 Case Processing.
- 6.2.13. Dispose of any remaining tissue. Refer to SOP 66 Specimen Disposal.

<b>JDRF nPOD Standard Operating Procedure</b>		
SOP Number and Version: 45.0	Supersedes: N/A	SPECIAL PROJECTS: ORGAN PROCUREMENT & PROCESSING CORE PROCEDURE
Original Effective Date: 08/27/2014		
Version Effective Date:		Page 6 of 7

## Procurement of Histological Specimens and Islets from The Same T1D Pancreas



JDRF nPOD Standard Operating Procedure			
SOP Number and Version:	45.0	Supersedes: N/A	SPECIAL PROJECTS: ORGAN PROCUREMENT & PROCESSING CORE PROCEDURE
Original Effective Date:	08/27/2014		
Version Effective Date:			
			Page 7 of 7